

A358 Taunton to Southfields Dualling Scheme

Ecological Baseline Report - Fish

HE551508-ARP-EBD-ZZ-RP-LE-000025

24 May 2022

Table of contents

	Pages
Executive summary	1
1 Introduction	2
1.1 Purpose and scope of this document	2
1.2 Scheme overview	2
1.3 Study area and zone of influence	3
1.4 Legislation	4
1.5 Status of notable fish species at the national level	4
1.6 Status of notable fish species at county level	5
1.7 Species-specific ecology for notable species	5
2 Methodology	7
2.1 Desk study	7
2.2 Field study	7
2.4 Fully-quantitative electric fishing survey	7
2.5 Qualitative electric fishing survey	8
2.6 Juvenile lamprey surveys	8
2.7 Habitat data and physico-chemical parameters	8
2.8 Biosecurity	8
2.9 Site selection	9
2.10 Assumptions and limitations	10
3 Results	11
3.1 Desk study	11
3.2 Field study	11
4 Conclusions	29
4.1 Survey constraints	29
Abbreviations List	31
Glossary	31
References	32
Appendices	i
Appendix A Fish survey study areas	ii
Appendix B EA desk study fish survey areas	iii
Appendix C Raw fish catch data	iv

Table of Figures

Figure 1-1	Scheme plan	3
Figure 3-1	Site 15a – Broughton Brook	12
Figure 3-2	Site 17 – Black Brook Tributary 1	12

Figure 3-3	Site 19a – Black Brook	13
Figure 3-4	Site 19b – Black Brook Tributary 3	13
Figure 3-5	Site 20 – Black Brook Tributary 2/5	14
Figure 3-6	Site 24 – Thornwater Stream	14
Figure 3-7	Site 30 – Meare Stream	15
Figure 3-8	Site 31 – Meare Stream Tributary 1	15
Figure 3-9	Downstream view at Fivehead River main channel 1	16
Figure 3-10	Site 34 species composition (actual catch)	17
Figure 3-11	Length-frequency histogram for minnow recorded at site 34	17
Figure 3-12	Length-frequency histogram for bullhead recorded at site 34	18
Figure 3-13	Length-frequency histogram for stoneloach recorded at site 34	18
Figure 3-14	Site 36 – Fivehead River main channel 2	19
Figure 3-15	Site 36 species composition (actual catch)	20
Figure 3-16	Length-frequency histogram for stoneloach recorded at site 36	20
Figure 3-17	Length-frequency histogram for 3-spined stickleback recorded at site 36	21
Figure 3-18	Site 37 – Fivehead River Tributary 5	21
Figure 3-19	Site 39 – Venner’s Water	22
Figure 3-20	Site 39 species composition (actual catch)	23
Figure 3-21	Site 51 – Cad Brook drainage network	23
Figure 3-22	Site 52 – Cad Brook	24
Figure 3-23	Site 54 -River Ding/River Ding drainage network	24
Figure 3-24	Site 54 species composition (actual catch)	25
Figure 3-25	Length-frequency histogram for 3-spined stickleback recorded at site 54	26
Figure 3-26	Length-frequency histogram for minnow recorded at site 54	26
Figure 3-27	Site 55 – Back Stream/River Ding drainage network	27
Figure 3-28	Site 55 species composition (actual catch)	28
Figure 3-29	Length-frequency histogram for minnow recorded at site 55	28
Figure 3-30	Length-frequency histogram for bullhead recorded at site 55	28

Table of Tables

Table 2-1	Summary table of fish survey locations	9
Table 2-2	Summary table of fish survey constraints	10
Table 3-1	Site 34 – In-situ water quality	16
Table 3-2	Site 34 catch summary	16
Table 3-3	Site 34 lamprey catch summary	18
Table 3-4	Site 36 – In-situ water quality	19
Table 3-5	Site 36 catch summary	20
Table 3-6	Site 39 – In-situ water quality	22
Table 3-7	Site 39 catch summary	22
Table 3-8	Site 54 – In-situ water quality	25
Table 3-9	Site 54 catch summary	25
Table 3-10	Site 55 – In-situ water quality	27
Table 3-11	Site 55 catch summary	27
Table 4-1	Summary table of fish survey constraints	29
Table 4-2	Summary table of notable fish species recorded at field survey sites	29
Table 4-3	Sites requiring additional survey	30
Table C-1	Species codes	v
Table C-2	Site 34 – Catch data	vi
Table C-3	Site 36 – Catch data	x
Table C-4	Site 39 – Catch data (qualitative)	xii
Table C-5	Site 54 – Catch data	xii

Table C-6 Site 55 – Catch data

Executive summary

The A358 Taunton to Southfields Dualling scheme would provide a dual carriageway along the length of the A358 between Taunton and Ilminster in Somerset, connecting the M5 motorway to the A303 to the south.

Fish surveys were part of the suite of habitat and protected species surveys commissioned in relation to the scheme. This report presents the results of the fish surveys undertaken throughout 2021 and aims to inform the ecology baseline for the scheme.

The objectives of this report are to present the methodologies used, identify survey limitations, and present the results of the fish surveys; the results of which will be used to inform appropriate mitigation and enhancement (if required).

A desk study undertaken reviewing available third-party data found the following records of fish species within the study area:

- Broughton Brook: brown trout (*Salmo trutta*), 3-spined stickleback (*Gasterosteus aculeatus*), European eel (*Anguilla anguilla*), bullhead (*Cottus gobio*) and stoneloach (*Barbatula barbatula*).
- Fivehead River: bullhead, minnow (*Phoxinus phoxinus*), stoneloach, brown trout, chub (*Squalius cephalus*), 3-spined stickleback, dace (*Leuciscus leuciscus*), European eel, and brook lamprey (*Lampetra planeri*).
- River Ding: European eel, bullhead, minnow and stoneloach.
- River Ilse: chub, dace, brown trout, minnow, bullhead, stoneloach, 3-spined stickleback and European eel.

Fish surveys were undertaken along a maximum 200 metre reach of any watercourses that were identified as being impacted by the scheme (100 metres upstream and downstream of the point the scheme intersects with the watercourse).

The majority of the watercourses within the study area were deemed unsuitable for electric fishing, due to either water depths/flow and/or the presence of dense vegetation/hedgerow inhibiting access and/or the application of electric fishing techniques.

At those sites which were suitable for electric fishing, a number of notable species were recorded including bullhead (Fivehead River Tributary 1 and Back Stream/River Ding drainage network), river lamprey (*Lampetra fluviatilis*)/brook lamprey (Fivehead River Tributary 1 and River Ding/River Ding drainage network) and brown trout (Fivehead River Tributary 1).

For those sites where the watercourse could not be seen clearly or where access was restricted due to the presence of dense vegetation, the need for further surveys when vegetation has died back (subject to avoiding fish spawning times and when water levels are suitable for surveying) has been identified.

For all sites where fish were encountered, mitigation is likely to be required where watercourses would be drained or diverted/realigned as part of scheme.

1 Introduction

1.1 Purpose and scope of this document

- 1.1.1 The A358 Taunton to Southfields Dualling scheme (hereafter referred to as ‘the scheme’) would provide a dual carriageway along the length of the A358 between Taunton and Ilminster in Somerset, connecting the M5 motorway to the A303. Electric fishing (fish) surveys were part of a suite of habitat and protected species surveys commissioned in relation to the scheme.
- 1.1.2 This report presents the results of the desk study and fish surveys and aims to inform the ecology baseline for the scheme.
- 1.1.3 The objectives of this report are to:
- summarise the results of the fish surveys, including analysis of the abundance, densities and age classes of each species
 - highlight the presence of any fish species of conservation interest
 - provide sufficient information to inform an assessment of the potential impacts to fish as a result of the scheme and design appropriate mitigation measures (where required)

1.2 Scheme overview

- 1.2.1 The scheme is part of a programme of improvements planned along the A303/A358 corridor aimed at improving connectivity between London, the south-east and the south-west. The A303, alongside the A30, forms part of the strategic road network (SRN) and together with the A358, provides the link between London, the south-east and the south-west.
- 1.2.2 The programme of improvements, as set out in the Government’s *Road Investment Strategy* [1] made a commitment to, “...upgrade all remaining sections of the A303 between the M3 and the A358 to dual carriageway standard, together with creating a dual carriageway link from M5 at Taunton to the A303”.
- 1.2.3 The scheme directly addresses this long-term commitment and would provide a new rural all-purpose dual carriageway link from the M5 at Taunton to the A303 at Southfields roundabout. The new dual carriageway would comprise new and upgraded stretches of the existing A358 road. Full details of the scheme will be provided in Chapter 2 *The Project* of the Environmental Statement (ES). Please refer to Figure 1-1 for a scheme plan.




Figure 1-1 Scheme plan

1.3 Study area and zone of influence

1.3.1 The Chartered Institute for Ecology and Environmental Management (CIEEM) *Guidelines for Ecological Impact Assessment* [2] recommend that all potentially important ecological features that occur within the Zone of Influence (Zoi) for a scheme are investigated. The Zoi includes:

- areas to be directly within the land take for the scheme
- areas that would be temporarily affected during construction
- areas likely to be impacted by hydrological disruption
- areas where there is a risk of pollution and noise disturbance during construction and/or operation

1.3.2 The Zoi depends on the ecological features concerned. With regard to the watercourses likely to be affected by the scheme in relation to fish, the Zoi has been defined as all watercourses that cross the scheme and all aquatic habitat 100 metres upstream and downstream of the crossing point, a total stretch of 200 metres of riverine habitat for each crossing point. This Zoi is hereafter referred to as the study area.

1.4 Legislation

1.4.1 A framework of international, European, national and local legislation and planning policy guidance exists to protect and conserve wildlife and habitats. This legislation will be listed in full within Chapter 8 *Biodiversity* of the ES. Legislation relevant to and discussed within this report are:

- The Conservation of Habitats and Species Regulations 2017 (the 'Habitats Regulations')
- Wildlife and Countryside Act 1981
- Natural Environment and Rural Communities (NERC) Act 2006
- Invasive Alien Species (Enforcement and Permitting) Order 2019
- UK Post-2010 Biodiversity Framework
- The Water Framework Directive (WFD) 2000/60/EC
- Salmon and Freshwater Fisheries (SAFFA) Act 1975
- Eels (England and Wales) Regulations 2009

1.4.2 It is an offence under the Salmon and Freshwater Fisheries Act 1975 to wilfully disturb spawning fish of any species, or habitat in which spawn is likely to be present. Any in-channel works should be timed to avoid spawning periods (March – June for coarse fish and November – January for salmonids). In addition, subject to subsections (3) and (5), any person who knowingly takes, kills or injures, or attempts to take, kill or injure, any salmon, trout, lamprey, smelt, shad, freshwater fish or specified fish in any waters which is unclean or immature shall be guilty of an offence.

1.5 Status of notable fish species at the national level

1.5.1 The status of notable fish species at the national level and details on species-specific ecology are presented below. Only species for which there were desk study or field survey records are discussed.

1.5.2 Bullhead (*Cottus gobio*): Bullhead is an EC Habitats Directive Annex II species. Core areas of habitats for species listed in Annex II are designated as sites of community importance (SCIs) and included in the Natura 2000 network.

1.5.3 River (*Lampetra fluviatilis*)/Brook (*Lampetra planeri*) lamprey: Lamprey are EC Habitats Directive Annex II species with core areas of habitats for these species listed in Annex II designated as SCIs and included in the Natura 2000 network. River lamprey is also listed as a species of principal importance (SPI) for the purpose of conserving biodiversity in accordance with the NERC Act 2006.

1.5.4 European eel (*Anguilla anguilla*): European eel is listed as Critically Endangered on the International Union for Conservation of Nature (IUCN) red list of threatened species. European eel is also listed as a SPI in accordance with the NERC Act 2006. The Eels (England and Wales) Regulations 2009 (Part 4) aims to maintain and improve the migratory routes of this species and requires specific mitigation around passage and screening in relation existing and proposed barriers to migration and abstractions.

1.5.5 Brown trout (*Salmo trutta*): Brown trout is listed as a SPI in accordance with the NERC Act 2006.

1.6 Status of notable fish species at county level

- 1.6.1 Although the UK Biodiversity Action Plan (UK BAP) has been superseded, BAPs are still widely used at county level to support the subsequent *UK Post-2010 Biodiversity Framework* [3]. Within *The Somerset Biodiversity Strategy 2008 – 2018* [4] water and wetlands are identified as a priority habitat, but individual fish species are not identified.

1.7 Species-specific ecology for notable species

Bullhead

- 1.7.1 Bullhead are a small bottom-dwelling fish with a preference for fast-flowing clear shallow water with a coarse substratum (gravel/cobble/pebble) [5]. Bullhead spawn from February to June, requiring a coarse substratum with large stones, providing habitat for the male to excavate a nest under a suitable large stone. Where large stones are not present, the bullhead may use woody debris or tree roots instead of large stones [6]. Newly hatched larvae are supported by a large yolk sac. Once the yolk has been exhausted, the fry are ready to disperse. Shallow, stony riffles are utilised by juvenile bullhead. Adult bullhead prefer sheltered sections created by woody debris, tree roots, leaf litter, macrophyte cover or large stones.
- 1.7.2 Generally, bullheads attain a length of 40–50mm after their first year, 60mm after their second and 70–90mm after their third. They do not generally live for more than three or four years [7].

Lamprey

- 1.7.3 Lamprey are a 'jawless' fish of the infraphylum *Agnatha* and are some of the most primitive vertebrates known to man. Three species of lamprey are found in the UK, these include: brook lamprey, river lamprey and sea lamprey (*Petromyzon marinus*). River and sea lamprey are anadromous species, taking to the sea as adults and returning to cool, clear headwater streams or clear gravels of main rivers to spawn. Brook lamprey do not feed during the adult phase of their lives and spawn shortly after metamorphosing into adults. Lamprey spawn on a stone/gravel substrate between March and June, where they create nests or redds [8]. Adult lamprey die shortly after breeding. Larvae emerge after 1-2 weeks and are known commonly as 'ammocoetes'.
- 1.7.4 Lamprey ammocoetes burrow themselves into the fine sand/silt, often found in river margins, where they feed by filtering micro-organisms such as algae, diatoms and detritus from the water column. After several years as ammocoete larvae, lamprey metamorphose into adult forms [9]. River and Sea lamprey migrate to estuaries and seas, where they adopt a hematophagous, parasitic, feeding strategy, before returning up-river to breed.

European eel

- 1.7.5 The European eel is a catadromous species, meaning that it migrates from freshwater to the sea to spawn. The breeding ground of the European eel is thought to be the Sargasso Sea. The larvae (leptocephalus), drift via the currents in the surface waters in the Atlantic, migrating to Europe [10]. Towards the end of this migration the larvae transform into a glass eel, which enter estuaries and migrate upstream. As the glass eels migrate upstream into freshwater

environments, they metamorphose into elvers. After five to 20 years in freshwater the eels reach sexual maturity, following which they begin their migration to their breeding grounds.

- 1.7.6 The European eel is largely nocturnal, resting under stones or burrowing into mud during the day, emerging at dusk in search of food [11]. Their diet is varied and includes marine, estuarine and freshwater fauna although they do not feed during the colder months [11]. The principal food items are invertebrates and small fish. Small eels feed on insect larvae, molluscs, worms and crustaceans while the diet of larger specimens consists predominately of other fish species.

Brown trout

- 1.7.7 Brown trout and sea trout are the same species but sea trout are anadromous, meaning they migrate to the sea to feed and return to freshwater to spawn [12]. Both sea and brown trout spawn upstream in freshwater where they lay their eggs in redds between November and January. The habitat requirements include gravel, good flow and a high level of oxygen. The eggs hatch into alevin which live in the gravel feeding off the remaining egg yolk. Once the yolk has been consumed, the alevin develop into fry. At this stage the trout will emerge from the gravel and feed upon insects towards the surface of the river.
- 1.7.8 Juvenile fish have a preference for particular substrate (e.g. gravel and cobble) and vegetation within which they seek refuge. Mature trout are larger but still vulnerable to predation and seek refuge along riverbanks, particularly those with submerged or overhanging tree roots, rocks or fallen logs. Adult trout require heterogeneous habitat types to provide a good food resource, whilst providing opportunities for shelter away from predator.

2 Methodology

2.1 Desk study

- 2.1.1 A combination of aerial photographs, Ordnance Survey maps and existing scheme information was used to identify watercourse scheme crossing points.
- 2.1.2 Information was obtained from the Somerset Environmental Records Centre (SERC) and the Environment Agency's Ecology & Fish Data Explorer [13] was used to identify the presence of any notable fish species records (collected within the past 10 years) from within a 2 kilometre radius or 1 kilometre upstream and downstream of where each watercourses is crossed by the scheme.
- 2.1.3 The results of the desk study are discussed by watercourse rather than by each individual crossing point.

2.2 Field study

- 2.2.1 Fish surveys were undertaken between 13 and 17 September 2021. This is within the optimal survey period, which is considered to be between August and early October following the end of the coarse fish spawning season (March – August) and prior to the salmonid spawning season (November – January).
- 2.2.2 The field surveys were led by Matthew Davidson, a Principal Aquatic Consultant and full member of the Institute of Fisheries Management (IFM) with 15 years' professional experience working within the commercial and academic sector.

2.3 Survey consent

- 2.3.1 Under section 27a of the Salmon and Freshwater Fisheries Act 1975, an FR2 'Application for authorisation to use fishing instruments other than rod and line in England' was submitted to the Environment Agency (EA), who provided a section 27a exception for the fish surveys. A copy of the Section 27a exception was in place prior to field survey operations.

2.4 Fully-quantitative electric fishing survey

- 2.4.1 Where possible, fully quantitative, catch depletion, electric-fishing surveys were performed using Electrafish backpack equipment. Survey reaches were chosen to encompass the range of available habitats to ensure a representative sample of the fish population was achieved. Where on-site conditions were suitable, a minimum of 50 metres survey length was sampled within the 200 metre study area.
- 2.4.2 Prior to surveys commencing at each site, fish populations were isolated using stop nets at the downstream and upstream survey extents. A minimum of three survey runs were undertaken within the netted reach. Studies have shown that three runs are usually sufficient to produce an accurate catch depletion analysis.
- 2.4.3 Sampling reaches were fished in an upstream direction and fish captured were netted and transferred to aerated holding containers for recovery and processing.
- 2.4.4 Once recovered, fish were identified and measured (fork length to the nearest millimetre). After processing, fish were returned to the same stretch of watercourse from which they were captured. Quantitative population density estimates for each fish species (number per m²) were calculated for each site,

based upon the depletion rate of consecutive catches taken from within the stop netted sections. Electric fishing was conducted by trained and highly experienced individuals following EA operational instruction guidance [14] and in accordance with BS EN 14011:2003 [15].

- 2.4.5 This approach enabled a population estimate for most sites/species to be made using the methodology provided by Carle & Strub (1978) [16]. This was calculated using the Pisces 2 'Population Estimation by Removal Sampling' software. In some instances, these calculations could not be made either due to insufficient numbers of captured fish or an irregular depletion (such as catching more fish in the second run than the first), in these instances population densities were calculated based upon the actual number of fish caught and as such can be viewed as a minimum estimate rather than a mathematical prediction of population densities.

2.5 Qualitative electric fishing survey

- 2.5.1 At site 39 (Venner's Water) a qualitative method was used to sample the fish population at this site. This method involved an electric fishing single run whereby fishing was undertaken by wading in an upstream direction as access allowed. This method enables a presence/absence assessment for the species present within the survey reach.

2.6 Juvenile lamprey surveys

- 2.6.1 Juvenile lamprey surveys, carried out between 13 and 17 September 2021, were undertaken at sites where suitable habitat was identified within or adjacent to the survey area. Suitable habitat for lamprey (fine sand or silt) was isolated with a net-walled quadrat and surveyed following standard methodology given in Natural England's publication *Monitoring the River, Brook and Sea Lamprey* [17]. A 1m² section was surveyed using on-off (20 seconds on, 5 seconds off) cycle draws with the anode positioned approximately 10 – 15cm above the substrate. For depletion purposes this process was repeated a minimum of three times with a 5-minute break between sampling rounds. Survey timings (between August and October), were considered to be optimal for larval lamprey surveys, ensuring a range of larval size classes and life stages would be captured [18].

2.7 Habitat data and physico-chemical parameters

- 2.7.1 A basic site description was completed including details such as the survey reach length, average width and depth and notable habitat features such as macrophyte stands, the presence of woody debris and the substrate composition. Basic physico-chemical parameters including temperature, pH, dissolved oxygen concentration, salinity and conductivity were also measured and recorded using a hand-held multi-parameter probe. Photographs of each survey reach were also taken to provide a visual record of the habitat surveyed and these are provided in Section 3.2.

2.8 Biosecurity

- 2.8.1 During the surveys surveyors adhered to strict biosecurity protocol to reduce the risk of transmitting invasive non-native species. Surveyors worked in an upstream to downstream direction on each watercourse. Before moving between

hydrologically unconnected watercourses, all equipment was thoroughly checked and cleaned.

2.9 Site selection

2.9.1 A total of 16 watercourse crossing locations were identified that interacted with the scheme (Table 2-1). Fish surveys were undertaken along a maximum 200 metre reach, 100 metres upstream and downstream of the point the scheme intersects with the watercourse (see Appendix A *Fish survey study areas*).

Table 2-1 Summary table of fish survey locations

Survey ID	Watercourse	Scheme crossing NGR	Survey methodology	Survey sample location	
				Upstream NGR	Downstream NGR
15a	Broughton Brook	ST 25593 24725	Survey not possible	-	-
17	Black Brook Tributary 1	ST 26005 24230	Survey not possible	-	-
19a	Black Brook	ST 25769 24572	Survey not possible	-	-
19b	Black Brook Tributary 3	ST 26525 23807	Survey not possible	-	-
20	Black Brook Tributary 2/5	ST 26306 23938	Survey not possible	-	-
24	Thornwater Stream	ST 27546 23555	Survey not possible	-	-
30	Meare Stream	ST 29245 21573	Survey not possible	-	-
31	Meare Stream Tributary 1	ST 29452 20806	Survey not possible	-	-
34	Fivehead River main channel 1	ST 29721 19292	50 metres fully quantitative and lamprey ammocoete survey	ST 29673 19246	ST 29641 19228
36	Fivehead River main channel 2	ST 30629 18516	50 metres fully quantitative	ST 30520 18474	ST 30552 18478
37	Fivehead River Tributary 5	ST 30827 18374	Survey not possible	-	-
39	Venner's Water	ST 31566 17915	10 metres qualitative	ST 31617 18020	-
51	Cad Brook drainage network	ST 33212 16561	Survey not possible	-	-
52	Cad Brook	ST 33224 10467	Survey not possible	-	-
54	River Ding/River Ding drainage network	ST 33609 15773	50 metres fully quantitative	ST 33632 15815	ST 33629 15794
55	Back Stream/River Ding drainage network	ST 33764 15708	50 metres fully quantitative	ST 33792 15715	ST 33826 15705

2.10 Assumptions and limitations

2.10.1 The survey results were affected by a number of issues as outlined below:

- Dense vegetation within and alongside the watercourse – dense vegetation coverage was encountered at the majority of the survey sites. The river channels were heavily vegetated and overgrown, primarily with rhizomatous and stoloniferous perennials and *Rubus* spp., amongst others; thus, making fully quantitative surveys impractical. The conditions encountered also presented an increased risk of injury or death to any fish populations from electric fishing due to the need for an increased number of runs to obtain a depletion and the likelihood of fish becoming entangled in the vegetation then being subjected to multiple runs (and therefore shocks) without capture.
- River hydrology – the majority of watercourses were observed to be slow flowing, ponded, and often dry and/or isolated in terms of their hydrology, therefore unsuitable for a fully quantitative survey. Aside from the conditions, if fish were present within those waterbodies the act of electric-fishing would put increased stress on them, potentially leading to mortality.

2.10.2 As a result of the constraints detailed above, it was only possible to undertake fully quantitative electric fishing surveys at four of the sixteen watercourse crossings, with a qualitative survey being undertaken at an additional site (site 39). Site locations and constraints are outlined in Table 2-2.

Table 2-2 Summary table of fish survey constraints

Survey ID	Watercourse	Constraint	Comment
34	Fivehead River main channel 1	substrate	Small species were difficult to catch due to nature of the substrate
36	Fivehead River main channel 2	flow, water turbidity & substrate	No perceptible flow upstream & downstream of survey reach. Questionable connectivity to main channel at time of survey. Highly turbid water and substrate composition reduced catch efficiency.
39	Venner's Water	water turbidity	Highly turbid water reduced catch efficiency
54	River Ding/River Ding drainage network	water depth & water turbidity	Deep, highly turbid water reduced catch efficiency.
55	Back Stream/River Ding drainage network	substrate & water turbidity	Small species were difficult to catch due to nature of the substrate. Highly turbid water reduced catch efficiency.

2.10.3 Where watercourses were listed as unsuitable for electric fishing, the assumption has been made that they are unlikely to support any notable fish species or significant fish population.

3 Results

3.1 Desk study

- 3.1.1 The SERC data search did not return any records of protected or notable fish species from within 2 kilometres of the study area.
- 3.1.2 The EA's Ecology & Fish Data Explorer [13] identified the presence of a number of notable fish species records, which are detailed below. The locations of the EA surveys are shown in Appendix B *EA desk study fish survey sites*.

Broughton Brook

- 3.1.3 A single run, semi-quantitative electric fishing survey of Broughton Brook was undertaken by the EA in March 2015 at New Barn Farm (ST 24997 23338) [13]. Analysis of the EA open data revealed the presence of brown trout, 3-spined stickleback (*Gasterosteus aculeatus*), European eel, bullhead and stoneloach (*Barbatula barbatula*).

Fivehead River

- 3.1.4 Two EA data sets were available for the Fivehead River, at Brandy Bridge (ST2900218966), to the south of the A358 and Elm Bridge (ST3139019670) to the north [13]. Electric fishing surveys were undertaken in July 2015 and April 2013. All sites had been completed using a single, semi quantitative electric fishing run.
- 3.1.5 The 2015 EA survey at Brandy Bridge revealed the presence of bullhead, minnow (*Phoxinus phoxinus*), stoneloach and brown trout. Elm Bridge was surveyed in 2010 and 2013 with the following species caught: chub (*Squalius cephalus*), 3-spined stickleback, minnow, dace (*Leuciscus leuciscus*), bullhead, European eel, brown trout, stoneloach and brook lamprey.

River Ding

- 3.1.6 The River Ding was surveyed by the EA in July 2015 at Broadway (ST3241715309) [13]. A single semi-quantitative electric fishing run was undertaken. The survey revealed the presence of European eel, bullhead, minnow and stoneloach.

River Isle

- 3.1.7 A semi-quantitative electric fishing survey undertaken by the EA at Hort Bridge (ST3464015326) in 2010 revealed the presence of chub, dace, brown trout, minnow, bullhead, stoneloach, 3-spined stickleback and European eels [13].

3.2 Field study

Site 15a – Broughton Brook

- 3.2.1 Site 15a was located to the east of the M5 junction 25, at the Nexus 25 roundabout, in an area of recent works associated with the new dual carriageway. The site was located under a pre-existing road crossing. The non-native invasive plant Himalayan balsam (*Impatiens glandulifera*) was extensive along both banks. The dense vegetation and hydrology made the site unsuitable for electric fishing within the study area.



Figure 3-1 Site 15a – Broughton Brook

Site 17 – Black Brook Tributary 1

- 3.2.2 Site 17 was a partly dry 'drainage ditch' with little to no flow and ponded areas at the time of survey. The channel was heavily vegetated with bankside vegetation consisting of broad-leaved trees, herbaceous plants and grasses, which shaded or completely covered the majority of the channel. The dense vegetation and hydrology made the site unsuitable for electric fishing within the study area.



Figure 3-2 Site 17 – Black Brook Tributary 1

Site 19a – Black Brook

- 3.2.3 Site 19a was located at the Nexus 25 roundabout, within an area of recent new development adjacent to the A358. The site was fenced and therefore inaccessible at the time of survey. The channel was heavily vegetated with bankside vegetation consisting of broad-leaved trees and herbaceous plants and grasses. River flow was minimal and consisted entirely of slack water, therefore conditions were unsuitable for electric fishing.



Figure 3-3 Site 19a – Black Brook

Site 19b – Black Brook Tributary 3

- 3.2.4 Site 19b was unsuitable for electric fishing. The river channel was completely covered with vegetation at the time of survey, consisting of dense hedgerow and herbaceous plants and grasses along both banks. Where possible to observe, water depth was shallow (<10cm) and ponded during the survey.



Figure 3-4 Site 19b – Black Brook Tributary 3

Site 20 – Black Brook Tributary 2/5

- 3.2.5 Site 20 was unsuitable for electric fishing. The site was heavily vegetated along both banks with dense hedgerow and low plants/grasses. The river channel was dry at the time of survey, both upstream and downstream of the scheme crossing point.



Figure 3-5 Site 20 – Black Brook Tributary 2/5

Site 24 – Thornwater Stream

- 3.2.6 Site 24 was unsuitable for electric fishing within the study area. Access to the river channel was restricted due to dense vegetation and coarse woody debris. Bankside vegetation was dominated by broad-leaved trees and herbaceous plants and grasses. The dense vegetation and hydrology made the site unsuitable for survey.



Figure 3-6 Site 24 – Thornwater Stream

Site 30 – Meare Stream

- 3.2.7 Site 30 was unsuitable for electric fishing within the study area. The river channel was bordered by dense bankside vegetation consisting of broad-leaved trees, dense hedgerow and herbaceous plants and grasses. The river channel was therefore inaccessible at the time of survey. Where observation allowed, water depth was shallow (<10cm), with little to no flow.



Figure 3-7 Site 30 – Meare Stream

Site 31 – Meare Stream Tributary 1

- 3.2.8 Site 31 was unsuitable for electric fishing. Water depths were shallow (<10cm) with little to no flow at the time of survey and the channel was heavily shaded. Bankside vegetation consisted of broad-leaved trees and herbaceous plants and grasses.



Figure 3-8 Site 31 – Meare Stream Tributary 1

Site 34 – Fivehead River main channel 1

- 3.2.9 The survey reach at site 34 was 50 metres in length and had an average width of approximately 5 metres. The approximate survey area was 250m². The maximum recorded depth was 0.4 metres with the average depth being approximately 0.25 metres. Dominant flow types included shallow run and glide, with some areas of slack water.



Figure 3-9 Downstream view at Fivehead River main channel 1

3.2.10 On the day of the survey the water was clear. The substrate consisted mainly of coarse sand/gravel, cobbles and fine sand/silt. Bankside vegetation consisted of broad-leaved trees, herbaceous plants and grasses. In-situ water quality readings are detailed in Table 3-1.

Table 3-1 Site 34 – In-situ water quality

Parameter	Reading
Temp °C	15.2
DO (%)	73.1
DO (mg/l)	7.35
pH	8.01
SPC (µs)	465.9
Sal (PPT)	0.23

3.2.11 A total of five species of fish were caught during the survey including minnow, stoneloach, 3-spined stickleback, brown trout and bullhead. The total number of individuals caught, catch depletion population estimates, densities and average (mean) lengths for each species are provided in Table 3-2 and further details are given in Appendix C *Raw fish data catch*. The community species composition is illustrated in Figure 3-10. Histograms showing the length-frequencies for minnow (Figure 3-10), bullhead (Figure 3-11) and stoneloach (Figure 3-12) are given below. Length-frequency histograms for the other species are not provided due to the low numbers of individuals caught.

Table 3-2 Site 34 catch summary

Species	Number caught	Catch depletion population estimate	Density per m ²	Fork length range (mm)	Average fork length (mm)
Minnow	50	57	0.228	18 - 85	53
Stoneloach	22	24	0.096	34 - 90	53
3-spined stickleback	1	n/a	0.004*	30 - 30	30
Brown trout	2	3	0.012	52 - 64	53

Species	Number caught	Catch depletion population estimate	Density per m ²	Fork length range (mm)	Average fork length (mm)
Bullhead	32	57	0.228	26 - 73	49

*minimum estimate




Figure 3-10 Site 34 species composition (actual catch)




Figure 3-11 Length-frequency histogram for minnow recorded at site 34




Figure 3-12 Length-frequency histogram for bullhead recorded at site 34




Figure 3-13 Length-frequency histogram for stoneloach recorded at site 34

- 3.2.12 A single juvenile lamprey (ammocoete) survey was undertaken at site 34, where a single 1m² quadrat of sub-optimal habitat was sampled. Ammocoetes were present and results are recorded in Table 3-3.
- 3.2.13 The habitat was sub-optimal, with substrate consisting of fine sand/silt. In this context, according to Harvey J & Cowx I (2003), optimal habitat is defined as “stable fine sediment or sand > 15 centimetres deep, low water velocity and the presence of organic detritus. Sub-optimal habitat is defined as shallow sediment, often patchy and interspersed among coarser substrate” [17]. Although it is recognised that the use of optimal/sub-optimal classification for lamprey habitat is no longer thought to be appropriate, as per recent guidance literature [18] it is considered a useful indicator for the purposes of this assessment in order to highlight the coarse substrate composition and therefore likely justification for the low numbers of lamprey spp. captured at the site.

Table 3-3 Site 34 lamprey catch summary

Run number	Species and life stage	Length (mm)		
		1	2	3
1	River/brook ammocoete	147	129	51
2	River/brook ammocoete	163	0	0
3	No catch	0	0	0

Site 36 – Fivehead River main channel 2

- 3.2.14 Site 36 was 50 metres in length and had an average width of approximately 1.5 metres. The total approximate survey area was 75m². The maximum recorded depth was 0.4 metres with the average depth being approximately 0.2 metres. Observed flow types were comprised entirely of shallow slack water.



Figure 3-14 Site 36 – Fivehead River main channel 2

- 3.2.15 On the day of the survey the water was slightly turbid, but the riverbed was visible throughout the survey reach. The substrate consisted mainly of coarse sand/gravel, with areas of fine sand/silt and cobbles/boulders. Bankside vegetation consisted of broad-leaved trees and herbaceous plants and grasses which shaded the channel. In-situ water quality readings are detailed in Table 3-4.

Table 3-4 Site 36 – In-situ water quality

Parameter	Reading
Temp °C	14.1
DO (%)	44.6
DO (mg/l)	10.3
pH	7.86
SPC (µs)	1054
Sal (PPT)	0.53

- 3.2.16 A total of three species of fish were caught during the survey at site 36: minnow, stoneloach, and 3-spined stickleback. The total number of individuals caught, catch depletion population estimates, densities and average (mean) lengths for each species are provided in Table 3-5 and further details are given in Appendix C *Raw fish data catch*. Species composition is illustrated in Figure 3-14 below. Histograms showing the length-frequencies for stoneloach and 3-spined stickleback are represented in Figure 3-15 and 3-16 respectively. A length-frequency histogram is not provided for minnow due to the low number of individuals caught.

Table 3-5 Site 36 catch summary

Species	Number caught	Catch depletion population estimate	Density per m ²	Fork length range (mm)	Average fork length (mm)
Minnow	1	n/a	0.013*	22 - 22	22
Stoneloach	6	7	0.093	102 - 132	67
3-spined stickleback	11	n/a	0.147*	25 - 58	43

*minimum estimate




Figure 3-15 Site 36 species composition (actual catch)




Figure 3-16 Length-frequency histogram for stoneloach recorded at site 36




Figure 3-17 Length-frequency histogram for 3-spined stickleback recorded at site 36

Site 37 – Fivehead River Tributary 5

- 3.2.17 Site 37 was a steep banked 'drainage ditch' unsuitable for electric fishing. The channel was heavily vegetated; bankside vegetation consisted of broad-leaved trees and herbaceous plants and grasses. Water depths were shallow (<10 centimetres) with little to no flow observed.



Figure 3-18 Site 37 – Fivehead River Tributary 5

Site 39 – Venner's Water

- 3.2.18 Access to the river channel was limited due to dense overhanging vegetation. A qualitative electric fishing survey was, however, undertaken along an approximately 10 metre reach of the river in order to establish presence/absence for fish species.
- 3.2.19 The survey reach at site 39 was approximately 10 metres in length and had an average width of 2 metres. The approximate survey area was 20m². The maximum recorded depth was 0.4 metres with the average depth being approximately 0.3 metres. Dominant flow types included shallow run and glide, with some areas of slack water.



Figure 3-19 Site 39 – Venner’s Water

3.2.20 On the day of the survey the water was highly turbid. The substrate consisted mainly of fine sand/silt, cobbles and coarse sand/gravel. Bankside vegetation consisted of broad-leaved trees, herbaceous plants and grasses and the river was heavily shaded. Himalayan balsam was also present along both banks. In-situ water quality readings are detailed in Table 3-6.

Table 3-6 Site 39 – In-situ water quality

Parameter	Reading
Temp °C	14.8
DO (%)	45.1
DO (mg/l)	4.45
pH	8.0
SPC (µs)	632
Sal (PPT)	0.31

3.2.21 A total of two species of fish were caught during the survey: stoneloach and 3-spined stickleback. The total number of individuals caught and average (mean) lengths for each species are provided in Table 3-7 and further details are given in Appendix C *Raw fish data catch*. Species composition is presented in Figure 3-19. Length-frequency histograms are not provided due to the low number of individuals caught.

Table 3-7 Site 39 catch summary

Species	Number caught	Fork length range (mm)	Average fork length (mm)
Stoneloach	1	115 - 115	115
3-spined stickleback	5	32 - 41	37




Figure 3-20 Site 39 species composition (actual catch)

Site 51 – Cad Brook drainage network

- 3.2.22 Site 51 was a heavily overgrown 'drainage ditch', unsuitable for electric fishing. There was no observable flow within the channel at the time of survey. Bankside vegetation consisted of dense hedgerow, bramble and herbaceous plants and grasses.



Figure 3-21 Site 51 – Cad Brook drainage network

Site 52 – Cad Brook

- 3.2.23 The river channel at site 52 was heavily vegetated at the time of survey; bankside vegetation consisted of herbaceous plants and grasses which obscured almost the entire channel. The dense vegetation and lack of open water made the site unsuitable for electric fishing at the time of survey.



Figure 3-22 Site 52 – Cad Brook

Site 54 – River Ding/River Ding drainage network

- 3.2.24 The survey reach at site 54 was 50 metres in length and had an average width of approximately 2 metres. The approximate survey area was 100m². The maximum recorded depth was 1.2 metres with the average depth being approximately 0.5 metres. Dominant flow types included deep run and glide, with some areas of marginal slack water.



Figure 3-23 Site 54 -River Ding/River Ding drainage network

- 3.2.25 On the day of the survey the water was highly turbid, which limited views of the substrate. Where substrate was visible, it consisted mainly of fine silt and clay. Bankside vegetation consisted of broad-leaved trees, herbaceous plants and grasses and the river was heavily shaded. In-situ water quality readings are detailed in Table 3-8.

Table 3-8 Site 54 – In-situ water quality


Parameter	Reading
Temp °C	15
DO (%)	50
DO (mg/l)	5.7
pH	7.64
SPC (µs)	423.6
Sal (PPT)	0.12

3.2.1 A total of three species of fish were caught during the survey: minnow, stone loach and 3-spined stickleback. The total number of individuals caught, catch depletion population estimates, densities and average (mean) lengths for each species are provided in Table 3-9 and further details are given in Appendix C *Raw fish data catch*. Species composition is illustrated in Figure 3-23. Histograms showing the length-frequencies for 3-spined stickleback (Figure 3-24) and minnow (Figure 3-25) are given below. A length-frequency histogram for stone loach is not provided due to the low numbers of individuals caught.

Table 3-9 Site 54 catch summary

Species	Number caught	Catch depletion population estimate	Density per m ²	Fork length range (mm)	Average fork length (mm)
Minnow	16	17	0.17	16 - 75	38
Stone loach	1	n/a	0.01*	47	87
3-spined stickleback	21	29	0.29	17 - 51	27

*minimum estimate

**Figure 3-24 Site 54 species composition (actual catch)**




Figure 3-25 Length-frequency histogram for 3-spined stickleback recorded at site 54




Figure 3-26 Length-frequency histogram for minnow recorded at site 54

3.2.2 A single, free-swimming sub-adult (macrophthalmia or transformer stage) brook lamprey was caught during the triple-shock survey. Following this, juvenile lamprey surveys were undertaken at the site, where a small area of suitable sub-optimal habitat was identified. No lamprey were caught.

Site 55 – Back Stream/River Ding drainage network

3.2.3 The survey reach at site 55 was 50 metres in length and had an average width of approximately 3 metres. The approximate survey area was 150m². The maximum recorded depth was 0.6 metres with the average depth being approximately 0.2 metres. Dominant flow types included shallow run and glide, with some areas of slack water and deeper glide and run.



Figure 3-27 Site 55 – Back Stream/River Ding drainage network

- 3.2.4 On the day of the survey the water was highly turbid, which limited views of the substrate. Where substrate was visible, it consisted mainly of coarse sand, gravel and cobbles with areas of fine sand/silt. Bankside vegetation consisted of broad-leaved trees, herbaceous plants and grasses and the river was heavily shaded by overhanging vegetation. In-situ water quality readings are detailed in Table 3-10.

Table 3-10 Site 55 – In-situ water quality

Parameter	Reading
Temp °C	15.7
DO (%)	70.7
DO (mg/l)	7.02
pH	7.86
SPC (µs)	430.9
Sal (PPT)	0.21

- 3.2.5 A total of four species of fish were caught during the survey: minnow, stoneloach, 3-spined stickleback and bullhead. The total number of individuals caught, catch depletion population estimates, densities and average (mean) lengths for each species are provided in Table 3-11 and further details are given in Appendix C *Raw fish data catch*. Species composition is illustrated in Figure 3-27. Histograms showing the length-frequencies for minnow (Figure 3-28) and bullhead (Figure 3-29) are given below. Length-frequency histograms for the other species are not provided due to the low numbers of individuals caught.

Table 3-11 Site 55 catch summary

Species	Number caught	Catch depletion population estimate	Density per m ²	Fork length range (mm)	Average fork length (mm)
Minnow	15	16	0.11	27-73	50
Bullhead	23	24	0.16	35 - 68	50
Stoneloach	1	n/a	0.01*	87	87
3-spined stickleback	1	n/a	0.01*	27	27

*minimum estimate




Figure 3-28 Site 55 species composition (actual catch)




Figure 3-29 Length-frequency histogram for minnow recorded at site 55




Figure 3-30 Length-frequency histogram for bullhead recorded at site 55

4 Conclusions

4.1 Survey constraints

4.1.1 Factors influencing the efficacy of surveys at each of the different sites are summarised in Table 4-1. Fully quantitative 50 metres surveys were undertaken at sites 34, 36, 54 and 55, whereas only a qualitative 10 metres survey was possible at site 39.

Table 4-1 Summary table of fish survey constraints

Survey ID	Watercourse	Constraint	Comment
34	Fivehead River main channel 1	substrate	Small species were difficult to catch due to nature of the substrate
36	Fivehead River main channel 2	flow, water turbidity & substrate	No perceptible flow upstream & downstream of survey reach. Questionable connectivity to main channel at time of survey. Highly turbid water and substrate composition reduced catch efficiency.
39	Venner's Water	water turbidity	Highly turbid water reduced catch efficiency
54	River Ding/River Ding drainage network	water depth & water turbidity	Deep, highly turbid water reduced catch efficiency.
55	Back Stream/River Ding drainage network	substrate & water turbidity	Small species were difficult to catch due to nature of the substrate. Highly turbid water reduced catch efficiency.

4.1.2 All the watercourses surveyed were small with narrow channel widths and predominantly shallow water depths. The fish species caught were typical for such watercourses, based on the habitat present. Although in some cases, specifically sites 36, 54 and 55, brown trout or eels could have been expected but were not encountered.

4.1.3 Species of note included bullhead, brown trout and river/brook lamprey ammocoete observed at the sites as summarised in Table 4-2.

Table 4-2 Summary table of notable fish species recorded at field survey sites

Survey ID	Watercourse	Notable species recorded during field survey
34	Fivehead River main channel 1	Bullhead, brown trout, river/brook lamprey ammocoete
36	Fivehead River main channel 2	N/A
39	Venner's Water	N/A
54	River Ding/River Ding drainage network	Brook lamprey (transformer)
55	Back Stream/River Ding drainage network	Bullhead

4.1.4 For those sites where the watercourse could not be seen clearly or where access was restricted due to the presence of dense vegetation, the need for further surveys when vegetation has died back (subject to avoiding fish spawning times and when water levels are suitable for surveying) has been identified. See Table 4-3 below for sites that have been identified as requiring further survey.

Table 4-3 Sites requiring additional survey

Survey ID	Watercourse	Survey timings*
15a	Broughton Brook	Spring
17	Black Brook Tributary 1	Spring
19a	Black Brook	Spring
19b	Black Brook Tributary 3	Spring
20	Black Brook Tributary 2/5	Spring
24	Thornwater Stream	Spring
30	Meare Stream	Spring
31	Meare Stream Tributary 1	Spring
37	Fivehead River Tributary 5	Spring
39	Venner's Water	Spring
51	Cad Brook drainage network	Spring
52	Cad Brook	Spring

* Subject to relevant EA consent

- 4.1.5 For all sites where fish were encountered mitigation is likely to be required if the watercourses are to be drained or diverted/realigned as part of the proposed work. Mitigation is likely to include a fish translocation and the presence of an aquatic ecological clerk of works whilst works are being undertaken.
- 4.1.6 As discussed in Section 1.5 of this report, it is an offence under the Salmon and Freshwater Fisheries Act 1975 to wilfully disturb spawning fish of any species, or habitat in which spawn is likely to be present. Any in-channel works should therefore be timed to avoid spawning periods (March – June for coarse fish and November – January for salmonids).

Abbreviations List

Please refer to ES Report Chapter 17 Abbreviations.

Glossary

Please refer to ES Report Chapter 18 Glossary.

References

- [1] Department for Transport, "Road investment strategy: 2015 to 2020," March 2015. [Online]. Available: <https://www.gov.uk/government/publications/road-investment-strategy-for-the-2010-to-2020-road-period>. [Accessed January 2021].
- [2] CIEEM, "Guidelines for Ecological Impact Assessment in the UK and Ireland: Terrestrial, freshwater, coastal and marine version 1.1," Chartered Institute for Ecology and Environmental Management, Winchester, 2018.
- [3] Joint Nature Conservation Committee, "UK Post-2010 Biodiversity Framework | JNCC - Adviser to Government on Nature Conservation," 2012. [Online]. Available: <https://jncc.gov.uk/our-work/uk-post-2010-biodiversity-framework/>. [Accessed October 2021].
- [4] Somerset Biodiversity Partnership, "Wild Somerset! 2008 - 2018. Somerset Biodiversity Strategy," [Online]. Available: <https://modgov.southsomerset.gov.uk/Data/South%20Somerset%20District%20Council/20090219/Agenda/Appendix%203%20-%20Somerset%20Biodiversity%20Strategy.pdf%2019-02-2009.pdf>. [Accessed October 2021].
- [5] W. J. P. Smyly, "The life-history of the bullhead or Miller's thumb (*Cottus gobio* L.)," *Proceedings of the Zoological Society of London*, vol. 128, no. 3, pp. 431-454, 1957.
- [6] N. T. Punchard, M. R. Perrow and A. J. D. Jowitt, "Fish Habitat Associations, Community Structure, Density and Biomass in Natural and Channelised Lowland Streams in the Catchment of the River Wensum, UK," in *Management and Ecology of River Fisheries*, 2000, p. Chapter 11.
- [7] P. Maitland and N. Campbell, *Freshwater Fishes of the British Isles*, London: Collins, 1992.
- [8] R. Gardiner and D. Stewart, "Spawning habitat assessment and survey of lamprey populations," Report to Scottish Natural Heritage. Contract Number RASD/060/96/N2K. Fisheries Research Services Report No. 07/97, Pitlochry, 1997.
- [9] M. W. Hardisty, "Information on the growth of the ammocoete larvae of the anadromous sea amprey, *Petromyzon marinus*, in British rivers," *Journal of Zoology*, vol. 159, no. 2, pp. 139-144, 1969.
- [10] A. Cresci, "A comprehensive hypothesis on the migration of European glass eels (*Anguilla anguilla*)," *Biological Reviews of the Cambridge Philosophical Society*, vol. 95, no. 5, pp. 1273-1286, 2020.
- [11] P. Walker, "Inside Ecology," Habitat preferences of the critically endangered European eel, December 2017. [Online]. Available: <https://insideecology.com/2017/12/22/habitat-preferences-of-the-critically-endangered-european-eel/>.

- [12] A. Ferguson, T. E. Reed, T. F. Cross, P. McGinnity and P. A. Prodohl, "Anadromy, potamodromy and residency in brown trout *Salmo trutta*: the role of genes and the environment," *Journal of Fish Biology*, pp. 1-27, 2019.
- [13] Environment Agency, "Ecology & Fish Data Explorer," 2021. [Online]. Available: <https://environment.data.gov.uk/ecology/explorer/>. [Accessed September 2021].
- [14] W. R. Beaumont, A. A. L. Taylor, M. J. Lee and J. S. Welton, "Guidelines for Electric Fishing Best Practice," *R&D Technical Report W2-054/TR*, 2002.
- [15] British Standards Institution, "S EN 14011:2003, BS 6068-5.32:2003: Water quality. Sampling of fish with electricity," 2003.
- [16] F. L. Carle and M. R. Strub, "A new method for estimating population size from removal data," *Biometrics*, vol. 34, pp. 621-630, 1978.
- [17] J. Harvey and I. Cowx, "Monitoring the River, Brook and Sea Lamprey, *Lampetra fluviatilis*, *L. planeri* and *Petromyzon marinus*. Conserving Natura 2000 Rivers Monitoring Series No. 5," English Nature, Peterborough, 2003.
- [18] JNCC, "Common Standards Monitoring Guidance for Freshwater Fauna, Version October 2015, ISSN 1743-8160," 2015. [Online]. Available: <https://data.jncc.gov.uk/data/9b80b827-b44b-4965-be8e-ff3b6cb39c8e/CSM-FreshwaterFauna-2015.pdf>.

Appendices

Appendix A Fish survey study areas




This map is reproduced from Ordnance Survey material with the permission of Ordnance Survey on behalf of the controller of Her Majesty's Stationery Office. © Crown Copyright. Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings. CLIENT NAME: Highways England LICENCE NUMBER: 100030649 [2021] Contains OS data © Crown Copyright and database right 2020 Source: Esri, Maxar, Earthstar Geographics, and the GIS User Community

LEGEND

- ECOLOGY SURVEY AREA
- ECOLOGY SURVEY AREA BUFFER - 100M
- WATERCOURSES
- FISH SURVEY EXTENT (100M EITHER SIDE OF SCHEME CROSSING POINT)

LABELS SHOW SURVEY LOCATION ID



SAFETY, HEALTH AND ENVIRONMENTAL INFORMATION	
IN ADDITION TO THE HAZARDS/RISKS NORMALLY ASSOCIATED WITH THE TYPES OF WORK DETAILED ON THIS DRAWING, NOTE THE FOLLOWING SIGNIFICANT RESIDUAL RISKS (REFERENCE SHALL ALSO BE MADE IN THE DESIGN HAZARD LOG)	
CONSTRUCTION	NONE
MAINTENANCE / CLEANING	NONE
USE	NONE
DECOMMISSIONING / DEMOLITION	NONE

Rev.	Date	Description	By	Chk'd	App'd	Auth'd
P02	03/05/22	ISSUE FOR INFORMATION	LL	MA	JS	SV

Suitability: S2
Drawing Status: SUITABLE FOR INFORMATION

TAYLOR WOODROW
together @ VINCI

ARUP **RAMBOLL**

national highways

Project Title: A358 TAUNTON TO SOUTHFIELDS				
Drawing Title: FISH SURVEY STUDY AREAS SHEET 1 OF 4				
Scale: 1:10,000	By: LL	Checked: MA	Approved: JS	Authorised: SV
Original Size: A3	Date: 03/05/22	Date: 03/05/22	Date: 03/05/22	Date: 03/05/22
Drawing Number: HE551508 - ZZ	HE PIN: ZZ	Originator: ARP	Volume: VES	Revision: P02
Location	Type	Role	Number	
	-DR-LE-000283			




This map is reproduced from Ordnance Survey material with the permission of Ordnance Survey on behalf of the controller of Her Majesty's Stationery Office. © Crown Copyright. Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings. CLIENT NAME: Highways England LICENCE NUMBER: 100030649 [2021] Contains OS data © Crown Copyright and database right 2020 Source: Esri, Maxar, Earthstar Geographics, and the GIS User Community

LEGEND

- ECOLOGY SURVEY AREA
- ECOLOGY SURVEY AREA BUFFER - 100M
- WATERCOURSES
- FISH SURVEY EXTENT (100M EITHER SIDE OF SCHEME CROSSING POINT)

LABELS SHOW SURVEY LOCATION ID



SAFETY, HEALTH AND ENVIRONMENTAL INFORMATION

IN ADDITION TO THE HAZARDS/RISKS NORMALLY ASSOCIATED WITH THE TYPES OF WORK DETAILED ON THIS DRAWING, NOTE THE FOLLOWING SIGNIFICANT RESIDUAL RISKS (REFERENCE SHALL ALSO BE MADE IN THE DESIGN HAZARD LOG)

CONSTRUCTION	NONE
MAINTENANCE / CLEANING	NONE
USE	NONE
DECOMMISSIONING / DEMOLITION	NONE

Rev.	Date	Description	By	Chk'd	App'd	Auth'd

Suitability: S2 | Drawing Status: SUITABLE FOR INFORMATION

TAYLOR WOODROW
together @ VINCI

ARUP | **RAMBOLL**

national highways

Project Title: A358 TAUNTON TO SOUTHFIELDS					
Drawing Title: FISH SURVEY STUDY AREAS SHEET 2 OF 4					
Scale: 1:10,000	By: LL	Checked: MA	Approved: JS	Authorised: SV	
Original Size: A3	Date: 03/05/22	Date: 03/05/22	Date: 03/05/22	Date: 03/05/22	
Drawing Number: HE551508 - ZZ	HE PIN: ZZ	Originator: ARP	Volume: VES	Revision: P02	
Location:	Type:	Role:	Number:	-DR - LE - 000284	




This map is reproduced from Ordnance Survey material with the permission of Ordnance Survey on behalf of the controller of Her Majesty's Stationery Office. © Crown Copyright. Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings. CLIENT NAME: Highways England LICENCE NUMBER: 100030649 [2021] Contains OS data © Crown Copyright and database right 2020 Source: Esri, Maxar, Earthstar Geographics, and the GIS User Community

LEGEND

- ECOLOGY SURVEY AREA
- ECOLOGY SURVEY AREA BUFFER - 100M
- WATERCOURSES
- FISH SURVEY EXTENT (100M EITHER SIDE OF SCHEME CROSSING POINT)

LABELS SHOW SURVEY LOCATION ID



SAFETY, HEALTH AND ENVIRONMENTAL INFORMATION	
IN ADDITION TO THE HAZARDS/RISKS NORMALLY ASSOCIATED WITH THE TYPES OF WORK DETAILED ON THIS DRAWING, NOTE THE FOLLOWING SIGNIFICANT RESIDUAL RISKS (REFERENCE SHALL ALSO BE MADE IN THE DESIGN HAZARD LOG)	
CONSTRUCTION	NONE
MAINTENANCE / CLEANING	NONE
USE	NONE
DECOMMISSIONING / DEMOLITION	NONE

Rev.	Date	Description	By	Chk'd	App'd	Auth'd
P02	03/05/22	ISSUE FOR INFORMATION	LL	MA	JS	SV


Suitability: S2 | Drawing Status: SUITABLE FOR INFORMATION

TAYLOR WOODROW
together @ VINCI

ARUP | **RAMBOLL**

national highways

Project Title		A358 TAUNTON TO SOUTHFIELDS			
Drawing Title		FISH SURVEY STUDY AREAS SHEET 3 OF 4			
Scale	By	Checked	Approved	Authorised	
1:10,000	LL	MA	JS	SV	
Original Size	Date	Date	Date	Date	
A3	03/05/22	03/05/22	03/05/22	03/05/22	
Drawing Number	HE PIN	Originator	Volume	Revision	
HE551508 - ZZ	ARP	- DR - LE - 000285	VES	P02	
Location	Type	Role	Number		




This map is reproduced from Ordnance Survey material with the permission of Ordnance Survey on behalf of the controller of Her Majesty's Stationery Office. © Crown Copyright. Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings. CLIENT NAME: Highways England LICENCE NUMBER: 100030649 [2021] Contains OS data © Crown Copyright and database right 2020 Source: Esri, Maxar, Earthstar Geographics, and the GIS User Community

LEGEND

- ECOLOGY SURVEY AREA
- ECOLOGY SURVEY AREA BUFFER - 100M
- WATERCOURSES
- FISH SURVEY EXTENT (100M EITHER SIDE OF SCHEME CROSSING POINT)

LABELS SHOW SURVEY LOCATION ID



SAFETY, HEALTH AND ENVIRONMENTAL INFORMATION	
IN ADDITION TO THE HAZARDS/RISKS NORMALLY ASSOCIATED WITH THE TYPES OF WORK DETAILED ON THIS DRAWING, NOTE THE FOLLOWING SIGNIFICANT RESIDUAL RISKS (REFERENCE SHALL ALSO BE MADE IN THE DESIGN HAZARD LOG)	
CONSTRUCTION	NONE
MAINTENANCE / CLEANING	NONE
USE	NONE
DECOMMISSIONING / DEMOLITION	NONE

Rev.	Date	Description	By	Chk'd	App'd	Auth'd
P02	03/05/22	ISSUE FOR INFORMATION	LL	MA	JS	SV

Suitability: S2 | Drawing Status: SUITABLE FOR INFORMATION

TAYLOR WOODROW
together @ VINCI

ARUP | **RAMBOLL**

national highways

Project Title		A358 TAUNTON TO SOUTHFIELDS			
Drawing Title		FISH SURVEY STUDY AREAS SHEET 4 OF 4			
Scale	By	Checked	Approved	Authorised	
1:10,000	LL	MA	JS	SV	
Original Size	Date	Date	Date	Date	
A3	03/05/22	03/05/22	03/05/22	03/05/22	
Drawing Number	HE PIN	Originator	Volume	Revision	
HE551508 - ZZ		ARP - DR - LE - 000286	VES	P02	
Location	Type	Role	Number		

Appendix B EA desk study fish survey areas



This map is reproduced from Ordnance Survey material with the permission of Ordnance Survey on behalf of the controller of Her Majesty's Stationery Office. © Crown Copyright. Unauthorised reproduction infringes Crown copyright and may lead to prosecution or civil proceedings. CLIENT NAME: Highways England LICENCE NUMBER: 100030649 [2021] Source: Esri, Maxar, GeoEye, Earthstar Geographics, CNES/Airbus DS, USDA, USGS, AeroGRID, IGN, and the GIS User Community

LEGEND	
	ECOLOGY SURVEY AREA
	ECOLOGY SURVEY AREA BUFFER - 100M
	EA DESK STUDY FISH SURVEY SITES

SAFETY, HEALTH AND ENVIRONMENTAL INFORMATION	
IN ADDITION TO THE HAZARDS/RISKS NORMALLY ASSOCIATED WITH THE TYPES OF WORK DETAILED ON THIS DRAWING, NOTE THE FOLLOWING SIGNIFICANT RESIDUAL RISKS (REFERENCE SHALL ALSO BE MADE IN THE DESIGN HAZARD LOG)	
CONSTRUCTION	
NONE	
MAINTENANCE / CLEANING	
NONE	
USE	
NONE	
DECOMMISSIONING / DEMOLITION	
NONE	

Rev.	Date	Description	By	Chk'd	App'd	Auth'd
P01	03/05/22	ISSUE FOR INFORMATION	LL	MA	JS	SV

S2 Drawing Status
 SUITABLE FOR INFORMATION

 together @ VINCI

 Client

Project Title		A358 TAUNTON TO SOUTHFIELDS	
Drawing Title			
EA DESK STUDY FISH SURVEY SITES			
Scale	1:50,000	By	LL
Original Size	A3	Date	03/05/22
Checked	MA	Date	03/05/22
Approved	JS	Date	03/05/22
Authorised	SV	Date	03/05/22
Drawing Number	HE551508 - ZZ	Originator	ARP - DR - LE - 000341
HE PIN	ZZ	Volume	VES
Location		Type	Role
		Number	Revision
			P01

Appendix C Raw fish catch data

C.1 Species Codes

Table C-1 Species codes

Code	Common name	Latin name
SP3	3-spined stickleback	<i>Gasterosteus aculeatus</i>
SP9	9-spined stickleback	<i>Pungitius pungitius</i>
SA	Atlantic salmon	<i>Salmo salar</i>
BA	Barbel	<i>Barbus barbus</i>
BI	Bitterling	<i>Rhodeus sericeus</i>
BL	Bleak	<i>Alburnus alburnus</i>
LB	Brook lamprey	<i>Lampetra planeri</i>
BT	Brown trout	<i>Salmo trutta</i>
BH	Bullhead	<i>Cottus gobio</i>
CC	Carp (Common)	<i>Cyprinus carpio (common)</i>
CR	Carp (Crucian)	<i>Carassius carassius</i>
GC	Carp (Ghost)	<i>Cyprinus carpio (ghost)</i>
KC	Carp (Koi)	<i>Cyprinus carpio (koi)</i>
LC	Carp (Leather)	<i>Cyprinus carpio (leather)</i>
MC	Carp (Mirror)	<i>Cyprinus carpio (mirror)</i>
GRC	Carp (Grass)	<i>Ctenopharyngodon idella</i>
GF	Goldfish	<i>Carassius auratus</i>
CH	Chub	<i>Squalius cephalus</i>
CB	Common bream	<i>Abramis brama</i>
CBXSB	Common bream x silver bream hybrid	<i>Abramis brama x Blicca bjoerkna</i>
DA	Dace	<i>Leuciscus leuciscus</i>
EE	European eel	<i>Anguilla anguilla</i>
FL	Flounder	<i>Platichthys flesus</i>
GR	Grayling	<i>Thymallus thymallus</i>
GU	Gudgeon	<i>Gobio gobio</i>
LSP	Lamprey spp	N/A
MI	Minnow	<i>Phoxinus phoxinus</i>
MM	Motherless minnow/sunbleak	<i>Leucaspis delineatus</i>
OR	Orfe varieties	<i>Leuciscus idus</i>
PE	Perch	<i>Perca fluviatilis</i>
PI	Pike	<i>Esox lucius</i>
RT	Rainbow trout	<i>Oncorhynchus mykiss</i>
LR	River lamprey	<i>Lampetra fluviatilis</i>
RO	Roach	<i>Rutilus rutilus</i>
ROXCB	Roach x common bream hybrid	<i>Rutilus rutilus x Abramis brama</i>
ROXCH	Roach x chub hybrid	<i>Rutilus rutilus x Leuciscus cephalus</i>

Code	Common name	Latin name
ROXRU	Roach x rudd hybrid	<i>Rutilus rutilus x Scardinius erythrophthalmus</i>
RF	Ruffe	<i>Gymnocephalus cernua</i>
RU	Rudd	<i>Scardinius erythrophthalmus</i>
LS	Sea lamprey	<i>Petromyzon marinus</i>
ST	Sea trout	<i>Salmo trutta</i>
SB	Silver bream	<i>Blicca bjoerkna</i>
SL	Stoneloach	<i>Barbatula barbatula</i>
TE	Tench	<i>Tinca tinca</i>
TMG	Top mouth gudgeon	<i>Pseudorasbora parva</i>
ZA	Zander	<i>Sander lucioperca</i>

C.2 Site 34

C.2.1 Fish Catch Data

Table C-2 Site 34 – Catch data

Run No.	Fish No.	Species	Length (mm)
1	1	MI	47
1	2	SL	86
1	3	MI	85
1	4	SL	70
1	5	MI	74
1	6	SL	67
1	7	MI	74
1	8	BT	64
1	9	SL	90
1	10	SL	79
1	11	MI	77
1	12	MI	73
1	13	SL	78
1	14	BH	60
1	15	SL	63
1	16	SL	74
1	17	SL	65
1	18	BH	62
1	19	BH	64
1	20	BH	67
1	21	BH	60
1	22	BH	29
1	23	MI	34
1	24	BH	68

Run No.	Fish No.	Species	Length (mm)
1	25	SL	63
1	26	SL	59
1	27	BH	48
1	28	BH	35
1	29	BH	35
1	30	MI	29
1	31	MI	40
1	32	MI	44
1	33	MI	52
1	34	MI	45
1	35	MI	47
1	36	MI	51
1	37	MI	44
1	38	MI	63
1	39	MI	23
1	40	MI	40
1	41	MI	44
1	42	MI	53
1	43	MI	40
1	44	MI	39
1	45	MI	25
1	46	MI	18
1	47	MI	20
1	48	MI	35
1	49	MI	24
1	50	MI	22
2	51	MI	26
2	52	BH	65
2	53	SL	81
2	54	MI	77
2	55	SL	77
2	56	SL	90
2	57	MI	27
2	58	BH	71
2	59	BH	61
2	60	MI	74
2	61	BT	52
2	62	BH	62
2	63	BH	65
2	64	BH	40
2	65	SL	77
2	66	MI	35

Run No.	Fish No.	Species	Length (mm)
2	67	MI	30
2	68	SL	64
2	69	MI	53
2	70	MI	50
2	71	SL	62
2	72	BH	49
2	73	MI	39
2	74	BH	56
2	75	SL	70
2	76	BH	30
2	77	BH	27
2	78	SP3	30
2	79	MI	28
2	80	BH	28
2	81	MI	39
2	82	MI	39
2	83	BH	40
2	84	BH	27
2	85	SL	34
3	86	MI	72
3	87	MI	79
3	88	MI	74
3	89	BH	73
3	90	MI	74
3	91	SL	70
3	92	MI	68
3	93	BH	66
3	94	BH	54
3	95	BH	50
3	96	BH	28
3	97	MI	35
3	98	MI	47
3	99	SL	59
3	100	MI	55
3	101	MI	28
3	102	SL	53
3	103	MI	33
3	104	BH	30
3	105	BH	26
3	106	BH	28
3	107	BH	28

C.2.2 Catch Depletion Analysis Statistics

Minnow

Result - Carle & Strub

Results

Estimated population = 57

Standard error = 4.829

Lower 95% confidence interval = 4.61387E6

Upper 95% confidence interval = 66.4651

Chi-squared = 1.33

Degrees of freedom = 1

Probability = 0.2485 Accept model ($P \geq 0.2$)

Number of observations = 50

Calculations

	Observed	Expected	Probability of capture
1	28	27.6699	0.485437
2	12	14.2379	0.485437

Stoneloach

Result - Carle & Strub

Results

Estimated population = 24

Standard error = 2.119

Lower 95% confidence interval = 4.61387E6

Upper 95% confidence interval = 28.1541

Chi-squared = 0.88

Degrees of freedom = 1

Probability = 0.3478 Accept model ($P \geq 0.2$)

Number of observations = 22

Calculations

	Observed	Expected	Probability of capture
1	11	12.5714	0.52381
2	8	5.98639	0.52381
3	3	2.85066	0.52381

Bullhead

Result - Carle & Strub

Results

Estimated population = 57
 Standard error = 19.008
 Lower 95% confidence interval = 4.61387E6
 Upper 95% confidence interval = 94.2549

Chi-squared = 1.77
 Degrees of freedom = 1
 Probability = 0.1838 Reject model (P<0.2)
 Number of observations = 32

Calculations

	Observed	Expected	Probability of capture
1	10	13.2174	0.231884
2	13	10.1525	0.231884
3	9	7.79829	0.231884

Brown Trout

Result - Carle & Strub

Results

Estimated population = 3
 Standard error = 2.739
 Lower 95% confidence interval = 4.61387E6
 Upper 95% confidence interval = 8.36768

Chi-squared = 0.57
 Degrees of freedom = 1
 Probability = 0.4486 Accept model (P>=0.2)
 Number of observations = 2

Calculations

	Observed	Expected	Probability of capture
1	1	1.2	0.4
2	1	0.72	0.4
3	0	0.432	0.4

C.3 Site 36

C.3.1 Fish Catch Data

Table C-3 Site 36 – Catch data

Run No.	Fish No.	Species	Length (mm)
1	1	SL	114
1	2	SL	132

Run No.	Fish No.	Species	Length (mm)
1	3	SL	105
1	4	SL	111
1	5	SP3	49
1	6	SP3	58
1	7	SL	102
1	8	MI	22
2	9	SP3	29
2	10	SP3	28
2	11	SP3	25
2	12	SP3	26
2	13	SP3	29
2	14	SL	110
3	15	SP3	28
3	16	SP3	28
3	17	SP3	27
3	18	SP3	40

C.3.2 Catch Depletion Analysis Statistics

Stoneloach

Result - Carle & Strub

Results

Estimated population = 7

Standard error = N/A error in method

Lower 95% confidence interval = N/A

Upper 95% confidence interval = N/A

Chi-squared = N/A

Degrees of freedom = N/A

Probability = N/A

Number of observations = 6

Calculations

	Observed	Expected	Probability of capture
1	5	4.66667	0.666667
2	1	1.55556	0.666667
3	0	0.518518	0.666667

C.4 Site 39

C.4.1 Fish Catch Data

Table C-4 Site 39 – Catch data (qualitative)

Run No.	Fish No.	Species	Length (mm)
1	1	SP3	32
1	2	SP3	38
1	3	SP3	38
1	4	SP3	41
1	5	SP3	36
1	6	SL	115

C.5 Site 54

C.5.1 Fish Catch Data

Table C-5 Site 54 – Catch data

Run No.	Fish No.	Species	Length (mm)
1	1	MI	27
1	2	MI	27
1	3	MI	35
1	4	SP3	30
1	5	SP3	47
1	6	SP3	45
1	7	SP3	30
1	8	SP3	42
1	9	SL	47
1	10	SP3	27
1	11	MI	18
1	12	MI	22
1	13	SP3	17
1	14	MI	20
1	15	MI	35
1	16	MI	42
1	17	MI	31
2	18	MI	75
2	19	MI	54
2	20	MI	32
2	21	SP3	29
2	22	SP3	26
2	23	SP3	31
2	24	SP3	49

Run No.	Fish No.	Species	Length (mm)
2	25	SP3	32
2	26	SP3	27
2	27	SP3	32
2	28	SP3	28
2	29	MI	17
2	30	MI	16
2	31	MI	18
2	32	SP3	24
2	33	LSP	187
3	34	MI	66
3	35	SP3	20
3	36	SP3	28
3	37	SP3	28
3	38	SP3	51
3	39	SP3	21

C.5.2 Catch Depletion Analysis Statistics

Minnow

Result - Carle & Strub

Results

Estimated population = 17

Standard error = 1.480

Lower 95% confidence interval = 4.61387E6

Upper 95% confidence interval = 19.9015

Chi-squared = 1.26

Degrees of freedom = 1

Probability = 0.2616 Accept model ($P \geq 0.2$)

Number of observations = 16

Calculations

	Observed	Expected	Probability of capture
1	9	10.0741	0.592593
2	6	4.10425	0.592593
3	1	1.6721	0.592593

3-spined stickleback

Result - Carle & Strub

Results

Estimated population = 29

Standard error = 6.767
 Lower 95% confidence interval = 4.61387E6
 Upper 95% confidence interval = 42.2627

Chi-squared = 1.84
 Degrees of freedom = 1
 Probability = 0.1746 Reject model (P<0.2)
 Number of observations = 21

Calculations

	Observed	Expected	Probability of capture
1	7	9.51563	0.328125
2	9	6.39331	0.328125
3	5	4.29551	0.328125

C.6 Site 55

C.6.1 Fish Data

Table C-6 Site 55 – Catch data

Run No.	Fish No.	Species	Length (mm)
1	1	MI	64
1	2	MI	53
1	3	BH	59
1	4	BH	58
1	5	BH	54
1	6	BH	56
1	7	BH	45
1	8	BH	47
1	9	SP3	27
1	10	BH	52
1	11	BH	50
1	12	BH	48
1	13	BH	40
1	14	BH	59
1	15	BH	44
1	16	BH	42
1	17	BH	52
1	18	BH	46
1	19	BH	38
1	20	MI	52
1	21	MI	38
1	22	MI	56

1	23	MI	60
1	24	MI	53
1	25	MI	56
1	26	BH	35
1	27	BH	46
1	28	BH	46
1	29	BH	40
1	30	BH	52
1	31	MI	50
1	32	MI	50
2	33	MI	54
2	34	MI	44
2	35	BH	47
2	36	MI	38
2	37	MI	27
2	38	SL	87
3	39	MI	73
3	40	BH	68

C.6.2 Catch Depletion Analysis Statistics

Minnow

Result - Carle & Strub

Results

Estimated population = 16

Standard error = 1.789

Lower 95% confidence interval = 4.61387E6

Upper 95% confidence interval = 19.5062

Chi-squared = 0.13

Degrees of freedom = 1

Probability = 0.7143 Accept model ($P \geq 0.2$)

Number of observations = 15

Calculations

	Observed	Expected	Probability of capture
1	10	10	0.625
2	4	3.75	0.625
3	1	1.40625	0.625

Bullhead

Result - Carle & Strub

Results

Estimated population = 24

Standard error = 3.025

Lower 95% confidence interval = 4.61387E6

Upper 95% confidence interval = 31.9294

Chi-squared = 0.77

Degrees of freedom = 1

Probability = 0.3810 Accept model ($P \geq 0.2$)

Number of observations = 23

Calculations

	Observed	Expected	Probability of capture
1	21	19.0345	0.793103
2	1	3.93817	0.793103
3	1	0.814793	0.793103

If you need help accessing this or any other National Highways information, please call **0300 123 5000** and we will help you.

© Crown copyright 2022.

You may re-use this information (not including logos) free of charge in any format or medium, under the terms of the Open Government Licence. To view this licence:

visit www.nationalarchives.gov.uk/doc/open-government-licence/

write to the **Information Policy Team, The National Archives, Kew, London TW9 4DU**, or email psi@nationalarchives.gsi.gov.uk.

Mapping (where present): © Crown copyright and database rights 2021 OS 100030649. You are permitted to use this data solely to enable you to respond to, or interact with, the organisation that provided you with the data. You are not permitted to copy, sub-licence, distribute or sell any of this data to third parties in any form.

This document is also available on our website at www.nationalhighways.co.uk

For an accessible version of this publication please call **0300 123 5000** and we will help you.

If you have any enquiries about this publication email info@nationalhighways.co.uk or call **0300 123 5000***. Please quote the National Highways publications code **PR91/22**.

*Calls to 03 numbers cost no more than a national rate call to an 01 or 02 number and must count towards any inclusive minutes in the same way as 01 and 02 calls.

These rules apply to calls from any type of line including mobile, BT, other fixed line or payphone. Calls may be recorded or monitored.

Printed on paper from well-managed forests and other controlled sources when issued directly by National Highways.

Registered office Bridge House, 1 Walnut Tree Close, Guildford GU1 4LZ

National Highways Company Limited registered in England and Wales number 09346363